

**Amendments to the Specification**

Please replace paragraph 3, beginning at line 27, with the following amended paragraph:

The observed inhibition of HSL by nicotinic acid treatment is mediated by a decrease in cellular cyclic adenosine monophosphate (cAMP) caused by the G-protein-mediated inhibition of adenylyl cyclase. Recently, the G-protein coupled receptors HM74 and HM74A have been identified as receptors for nicotinic acid (PCT patent application WO02/84298; Wise et. al. J Biol Chem. 2003 **278** (11) 9869-9874). The DNA sequence of human HM74A (SEQ ID NOS: 1 and 2) may be found in Genbank; accession number AY148884. Two other papers support this discovery, (Tunaru et. al. Nature Medicine 2003 (3) 352-255 and Soga et. al. Biochem Biophys Res Commun. 2003 **303** (1) 364-369), however the nomenclature differs slightly. In the Tunaru paper what they term human HM74 is in fact HM74A and in the Soga paper HM74b is identical to HM74A. Cells transfected to express HM74A and/or HM74 gain the ability to elicit G<sub>i</sub> G-protein mediated responses following exposure to nicotinic acid. In mice lacking the homologue of HM74A (m-PUMA-G) nicotinic acid fails to reduce plasma NEFA levels.

Please insert the attached paper copy of the Sequence Listing into the Specification.